

Path Integral Over Conformally Self-Dual Geometries[†]Christof Schmidhuber^{*}*California Institute of Technology, Pasadena, CA 91125***Abstract**

The path integral of four dimensional quantum gravity is restricted to conformally self-dual metrics. It reduces to integrals over the conformal factor and over the moduli space of conformally self-dual metrics and can be studied with the methods of two dimensional quantum gravity in conformal gauge. The conformal anomaly induces an analog of the Liouville action. The proposal of David, Distler and Kawai is generalized to four dimensions. Critical exponents and the analog of the $c = 1$ barrier of two dimensional gravity are derived. Connections with Weyl gravity and four dimensional topological gravity are suggested.

hep-th@xxx/9112005

[†] Work supported in part by the U.S. Department of Energy under Contract No. DE-AC 0381-ER40050.

^{*} christof@theory3.caltech.edu

1. Introduction

The path integral of two dimensional quantum gravity in conformal gauge essentially reduces to an integral over the conformal factor. Its dynamics are provided by the conformal anomaly which can be integrated to give the Liouville action.^[1] David, Distler and Kawai (DDK) proposed reformulating the theory in terms of free fields propagating in a fictitious gravitational background with a marginal operator in the role of the cosmological constant term.^{[2][3]} The scaling laws derived from this proposal, e.g. for the fixed area partition function $Z(A)$ at zero cosmological constant, are in agreement with the results from light cone gauge,^[4] computer simulations^[5] and matrix models.^[6] These laws have been derived for two dimensional quantum gravity coupled to conformal field theories with $c \leq 1$, including nonunitary ones.

It is natural to ask whether we can apply the same methods to four dimensional quantum gravity and, e.g., derive the analog of the $c = 1$ barrier or scaling laws that could be compared with computer simulations. In this paper I show that this is the case at least if we restrict ourselves to conformally self-dual metrics: the path integral to study is

$$\int Dg Dx Dp e^{-S_{mat}[g,x] - \int_M d^4x \sqrt{g}(\lambda + \gamma R + \eta R^2 + ipW_+)} O_1[g,x] \dots O_n[g,x] \quad (1.1)$$

with a Lagrange multiplier p . The manifold M has fixed topology and Euclidean signature. x represents some conformally invariant matter fields, $O_1 \dots O_n$ are local operators, λ and γ are the cosmological and inverse Newtonian constants. The R^2 -term will be needed to cancel an R^2 -term from the conformal anomaly and will disappear in the end. $W_{\pm}$ is the (anti-) self-dual part of the (traceless) Weyl tensor

$$W_{\pm \mu\nu\sigma\tau} \equiv \frac{1}{2}(W_{\mu\nu\sigma\tau} \pm \frac{1}{2}\epsilon_{\mu\nu}^{\alpha\beta}W_{\alpha\beta\sigma\tau}),$$

$$W_{\mu\nu\sigma\tau} = R_{\mu\nu\sigma\tau} - \frac{1}{2}(g_{\mu\sigma}R_{\nu\tau} + g_{\nu\tau}R_{\mu\sigma} - g_{\mu\tau}R_{\nu\sigma} - g_{\nu\sigma}R_{\mu\tau}) + \frac{1}{6}(g_{\mu\sigma}g_{\nu\tau} - g_{\mu\tau}g_{\nu\sigma})R.$$

The Lagrange multiplier p is a 4th rank self-dual tensor field which (like W_+) transforms as (2,0) under the Euclideanized Lorentz group $SO(4) \sim SU(2) \times SU(2)$, i.e., like a spin 2 field.

The merit of p is that it reduces the path integral over the metric to an integral over the finite dimensional moduli space of conformally (anti-) self-dual metrics, the conformal factor and the diffeomorphism group, as will be explained. This makes (1.1) similar to two dimensional quantum gravity, the moduli space of conformally self-dual metrics playing the role of the moduli space of Riemann surfaces. As a consequence, we will be able to apply the same methods.

Why is “conformally (anti-) self-dual quantum gravity” interesting? One reason is that it is a four dimensional model for gravity that we might be able to quantize consistently and solve and in which we can nicely study, e.g., the effects of the conformal anomaly. It is a much richer toy model than two dimensional quantum gravity, because there the one dimensional universes have no geometry, only a total length, and the wave function of the universe is therefore relatively trivial. Conformally self-dual gravity will also be argued to describe renormalization group fixed points of gravity with a Weyl term, characterized by infinite Weyl coupling constant. One might speculate that they correspond to long or short distance “phases”. One might also hope that some of the results found in this infinite coupling limit prevail qualitatively at finite or zero Weyl coupling.

Of course, there is a well known ghost problem common to all fourth order derivative actions (like the Weyl action:) we can rewrite them in terms of new fields with two derivatives only, but some of them will have the wrong sign in the kinetic term. With Minkowskian signature this leads to nonunitarity. Two types of fourth order derivative terms will arise in this paper: terms quadratic in the curvature, and a new term induced by the conformal anomaly which makes the model renormalizable. The former will either be tuned away in the end or decouple. The ghosts arising from the latter might decouple due to the reparametrization constraints, as in string theory. In any case, they will not affect the calculations done.

Part of this paper is concerned with the four dimensional analog of the Liouville action and of DDK. In a different context the induced action for the conformal factor and its renormalization have also been studied recently by Antoniadis and Mottola. I

have used some of their calculations.^[7] However, when I discuss the four dimensional analog of DDK's method of decoupling the conformal factor from its measure, my treatment and my conclusions will differ from those of [7]. I will state the main differences.

In section two, (1.1) without the operators will be rewritten as an integral over moduli space and over the conformal factor ϕ with a few determinants in its gravitational background. As in two dimensions, the determinants can be decoupled from ϕ by introducing a 4D analog of the Liouville action. Its form has already been found in [7]. It consists of a free 4th order derivative piece (essentially $\phi \square^2 \phi$) plus pieces that renormalize λ, γ and η in (1.1), as explained in section 3.

In quantum gravity the measure $D_g g$ for the metric is defined with respect to the fluctuating metric itself. In two dimensions, DDK (further elaborated in^[8]) replaced the conformal factor by a field whose measure is defined with respect to some background metric and whose action is again the Liouville action with modified coefficients, determined by requiring that the choice for the background metric is irrelevant. In chapter 4, the same is done in four dimensions. The cosmological constant, the Hilbert-Einstein term and the R^2 term should each become truly marginal operators of the new theory, but so far I have verified this only for the cosmological term.

In section 5 scaling laws in conformally self-dual quantum gravity are derived, similar to the two dimensional ones.* I will discuss the partition function at fixed volume or average curvature, and the correlation functions of local operators in their dependence on the cosmological constant. It would be very interesting to explore whether the condition $W_+ = 0$ can be imposed in computer simulations of random triangulations. Then these predictions could be compared with "experiment."

Section 6 discusses the connection of (1.1) with fixed points of gravity with a Weyl term. It is also suggested that conformally self-dual gravity is connected with four dimensional topological gravity,^[9] as in the two dimensional case.^[10]

* the values of the exponents and the analog of the $c = 1$ barrier can be found in the note added at the end of the paper

2. Conformal Gauge

The Lagrange multiplier p in (1.1) restricts the path integral over g to conformally self-dual metrics, i.e. metrics with $W_+ = 0$. W_+ has five independent components and the condition $W_+ = 0$ is Weyl- and diffeomorphism invariant. So up to a finite number of moduli the five surviving components of the metric will be the conformal factor and the diffeomorphisms. Let m_i parametrize the moduli space of conformally self-dual metrics modulo diffeomorphisms $x \rightarrow x + \xi$ and Weyl transformations $g \rightarrow ge^\phi$. Let us fix a representative $\hat{g}(m_i)$ via, say, the condition $\hat{R} = 0$ and Lorentz gauge $\partial^\mu \hat{g}_{\mu\nu} = 0$ and let us pick a conformally self dual metric

$$g_0 = (\hat{g}(m_i)e^\phi)^\xi \quad (2.1)$$

where ξ indicates the action of a diffeomorphism. At g_0 we can split up g :

$$\delta g_{\mu\nu} = g_{0\mu\nu} \delta\phi + \nabla_{(\mu} \delta\xi_{\nu)} + \delta\bar{h}_{\mu\nu}.$$

The four $\delta\xi$'s generate infinitesimal diffeomorphisms and the five $\bar{h}_{\mu\nu}$ parametrize the space of metrics perpendicular to ξ , ϕ and the moduli, i.e., perpendicular to the conformally self dual ones. The measure for g is defined, in analogy to two dimensions^[1], by

$$\|\delta g\|^2 \equiv \int d^4x \sqrt{g} (4(\delta\phi + \frac{1}{2}\nabla^\mu \delta\xi_\mu)^2 + (L\delta\xi)^2 + (\delta\bar{h})^2) \quad (2.2)$$

with

$$(L\delta\xi)_{\mu\nu} \equiv \nabla_{(\mu} \delta\xi_{\nu)} - \frac{1}{2}g_{\mu\nu} \nabla^\rho \delta\xi_\rho. \quad (2.3)$$

Apart from restricting the path integral, integrating out p and $\bar{h}$ in (1.1) will contribute the determinant

$$\det(O^\dagger O)_{g_0}^{-\frac{1}{2}} \quad (2.4)$$

where $O^\dagger$ is the linearized W_+ -term

$$(O_{g_0}^\dagger \bar{h})_{\mu\nu\sigma\tau} \equiv \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} (W_{+ \mu\nu\sigma\tau}[g_0 + \epsilon \bar{h}] - W_{+ \mu\nu\sigma\tau}[g_0]), \quad (2.5)$$

O is its adjoint and $O^\dagger O$ is a 4^{th} order, conformally invariant, linear differential operator in the curved background g_0 , acting on p .

We are left with an integral over the conformal equivalence class of each $\hat{g}$. From (2.2) it is seen that changing variables from g to ϕ and ξ in this equivalence class leads to a Jacobian

$$\det(L^\dagger L)_g^{\frac{1}{2}}$$

where the zero modes of the operator L , defined in (2.3), have to be projected out. After dropping the integral over the diffeomorphism group $D\xi$ (since gravitational anomalies can occur only in $4k+2$ dimensions^[11]) the path integral (1.1) without the operator insertions reduces to an integral over the moduli space of conformally self-dual metrics and ϕ :

$$\int \prod_i dm_i D\phi \det(O^\dagger O)_{\hat{g}e^\phi}^{-\frac{1}{2}} \det(L^\dagger L)_{\hat{g}e^\phi}^{\frac{1}{2}} \det(\Delta)_{\hat{g}e^\phi}^{-\frac{1}{2}} e^{-\int d^4x \sqrt{g}(\lambda + \gamma R + \eta R^2)} \quad (2.6)$$

where the matter partition function has been denoted by $\det(\Delta)^{-\frac{1}{2}}$. Despite of the notation, the conformally invariant matter is allowed to be fermions, Yang-Mills fields, etc., as well as conformally coupled scalars.

The moduli space of conformally self-dual metrics is a very interesting subject by itself which will not be discussed here. On the four sphere its dimension is zero: all conformally self-dual metrics on S^4 are conformally flat. On K^3 , e.g., its dimension is 57.^[12]

3. Liouville in 4D

Let us now decouple the determinants in (2.6) from ϕ . For conformally invariant differential operators X :

$$\det X_{\hat{g}e^\phi} = \det X_{\hat{g}} e^{-S_i[\hat{g}, \phi]} \quad (3.1)$$

where the induced action S_i is obtained from integrating the trace anomaly of the stress tensor^[13]

$$-2 \frac{\delta S_i[\hat{g}, \phi]}{\delta \phi} = \sqrt{g} \langle T^\mu_\mu \rangle = \frac{1}{16\pi^2} \sqrt{g} [a(F + \frac{2}{3} \square R) + bG] - 4\lambda' \sqrt{g} - 2\gamma' \sqrt{g} R \quad (3.2)$$

where $F = W_+^2 + W_-^2$ is the square of the Weyl tensor. (3.2) has, apart from the divergent parameters λ' and γ' , two finite parameters a, b . $\sqrt{g}G$ is the Gauss-Bonnet density whose integral over the manifold is proportional to the Euler characteristic. Following Antoniadis and Mottola^[7], (3.2) can actually easily be integrated by noting that with $g = \hat{g}e^\phi$ the combination

$$\sqrt{g}(G - \frac{2}{3} \square R) = \sqrt{\hat{g}} \hat{M} \phi + \sqrt{\hat{g}} (\hat{G} - \frac{2}{3} \square \hat{R})$$

is only linear in ϕ with the fourth order differential operator

$$\begin{aligned} \hat{M} &\equiv 2 \hat{\square}^2 + 4\hat{R}^{\mu\nu} \hat{\nabla}_\mu \hat{\nabla}_\nu - \frac{4}{3} \hat{R} \hat{\square} + \frac{2}{3} (\hat{\nabla}^\mu \hat{R}) \hat{\nabla}_\mu \\ &= 2 \hat{\square}^2 + 4\hat{R}^{\mu\nu} \hat{\nabla}_\mu \hat{\nabla}_\nu \quad \text{if } \hat{R} = 0 \\ &= 2 \hat{\square}^2 \quad \text{if } \hat{g} = \delta e^{\phi_0}. \end{aligned} \quad (3.3)$$

$\sqrt{g}F$ is independent of ϕ and $\sqrt{g} \square R$ integrates to the R^2 action. So the four dimensional analog of the Liouville action consists of a free part plus a cosmological constant term, a Hilbert-Einstein term and an R^2 term:

* More precisely, if $X \equiv M^\dagger M$, M has to transform as $e^{p\phi} M e^{q\phi}$ under $g \rightarrow ge^\phi$. It can be shown that L transforms as $e^\phi L e^{-\phi}$.

$$S_i[\hat{g}, \phi] = \frac{-b}{32\pi^2} S_0[\hat{g}, \phi] + \frac{-a}{32\pi^2} S_1[\hat{g}, \phi] + \frac{a+b}{72\pi^2} S_{R^2} + \gamma' S_R + \lambda' S_{c.c.} \quad (3.4)$$

where

$$\begin{aligned} S_0[\hat{g}, \phi] &= \int d^4x \sqrt{\hat{g}} \left[\frac{1}{2} \phi \hat{M} \phi + \left(\hat{G} - \frac{2}{3} \hat{\square} \hat{R} \right) \phi \right] \\ S_1[\hat{g}, \phi] &= \int d^4x \sqrt{\hat{g}} \hat{F} \phi \\ S_{c.c.} &= \int d^4x \sqrt{\hat{g}} e^{2\phi} \\ S_R &= \int d^4x \sqrt{\hat{g}} e^\phi \left[\hat{R} - \frac{3}{2} (\hat{\nabla} \phi)^2 - 3 \hat{\square} \phi \right] \\ S_{R^2} &= \int d^4x \sqrt{\hat{g}} \left[\hat{R} - \frac{3}{2} (\hat{\nabla} \phi)^2 - 3 \hat{\square} \phi \right]^2 \end{aligned} \quad (3.5)$$

b will turn out to be negative for ‘normal’ operators X . γ', λ' and $\frac{a+b}{72\pi^2}$ just renormalize γ, λ and η . A ϕ -independent local term

$$- \int d^4x \sqrt{\hat{g}} \left(\frac{a+b}{72\pi^2} \hat{R}^2 + \gamma' \hat{R} + \lambda' \right) \quad (3.6)$$

has been omitted in (3.4) and will frequently be omitted in the following. If it is included we see from (3.1) that for some action S_j :

$$S_i[\hat{g}, \phi] = S_j[\hat{g} e^\phi] - S_j[\hat{g}]. \quad (3.7)$$

$\frac{-b}{32\pi^2} S_0$ is the 4D analog of the 2D action

$$S_{2D} = \frac{c}{48\pi} \int d^4x \sqrt{\hat{g}} \left(\frac{1}{2} \phi \hat{\square} \phi - \hat{R} \phi \right).$$

If $\hat{g} = \tilde{g} e^{\phi_0}$, S_0 can be written:

$$S_0[\hat{g}, \phi] = \int d^4x \sqrt{\tilde{g}} \frac{1}{2} [(\phi + \phi_0) \tilde{M}(\phi + \phi_0) - \phi_0 \tilde{M} \phi_0]. \quad (3.8)$$

Adding up the anomaly coefficients in (3.2) for $\det(O^\dagger O)^{-\frac{1}{2}}, \det(L^\dagger L)^{+\frac{1}{2}}, \det \Delta^{-\frac{1}{2}}$,

$$A_0 \equiv a_O + a_L + a_{mat} \quad B_0 \equiv b_O + b_L + b_{mat}, \quad (3.9)$$

(2.6) can now be rewritten as

$$\int \prod_i dm_i \chi(m_i) \int D\phi e^{\frac{B_0}{32\pi^2} S_0[\hat{g}, \phi] + \frac{A_0}{32\pi^2} [\hat{g}, \phi] - \eta_1 S_{R^2} - \gamma_1 S_R - \lambda_1 S_{c.c.}} \quad (3.10)$$

$$\chi(m_i) \equiv \det(O^\dagger O)_{\hat{g}(m_i)}^{-\frac{1}{2}} \det(L^\dagger L)_{\hat{g}(m_i)}^{\frac{1}{2}} \det(\Delta)_{\hat{g}(m_i)}^{-\frac{1}{2}}$$

$\chi(m_i)$ is now purely a function of the moduli m_i , once we have fixed a representative $\hat{g}(m_i)$ for each point in moduli space.

The coefficients a and b in (3.2) have been calculated for, e.g., ^[13] conformally coupled scalars ($\Delta \sim \square - \frac{1}{6}R$):

$a_0 = \frac{1}{120}$	$b_0 = -\frac{1}{360}$
spin $\frac{1}{2}$ (four component) fermions:	$a_{\frac{1}{2}} = \frac{6}{120}$ $b_{\frac{1}{2}} = -\frac{11}{360}$
massless gauge fields:	$a_1 = \frac{12}{120}$ $b_1 = -\frac{62}{360}$

(3.11)

a_O, a_L, b_O, b_L , as well as a_M, b_M of M in (3.3), which will also be needed later, can in principle be calculated as usual with the Schwinger-de Witt method.* This will be tedious, especially for the fourth order derivative operator $O^\dagger O$ which acts on a 4th rank tensor field and is the sum of many components. It may be easier to study the operator product expansion of the *flat space* stress tensor with itself and to see whether one can infer the anomaly coefficients a, b from it, as in two dimensions. There the central charge c is read off from

$$T(z)T(w) \sim \frac{\frac{c}{2}}{(z-w)^2} + \dots$$

Note that the fourth order derivative induced action makes the theory power counting renormalizable, and also bounded if $b < 0$. The price is the existence of a ghost, the general problem of fourth order derivative actions mentioned in the introduction. It has already been suggested in [7] that the reparametrization constraints $T_{\mu\nu} \sim 0$ eliminate these ghosts from the physical spectrum as they do in two dimensions.^[14] This will be very interesting to explore in the future.

* see the note added at the end of the paper for the values.

4. David, Distler and Kawai in 4D

Let us now focus on the ϕ integral over the conformal equivalence class of $\hat{g}$:

$$Z[\hat{g}] \equiv \int D_{\hat{g}e^\phi} \phi e^{\frac{B_0}{32\pi^2}S_0[\hat{g},\phi] + \frac{A_0}{32\pi^2}S_1[\hat{g},\phi] - \eta_1 S_{R^2} - \gamma_1 S_R - \lambda_1 S_{c.c.}} \quad (4.1)$$

where the dependence of Z on η_1, γ_1 and λ_1 has been suppressed. In $D_{\hat{g}e^\phi}\phi$ it is indicated that the measure for ϕ depends on ϕ itself, namely in two ways: First, the metric itself must be used to define a norm in the space of metrics:

$$\|\delta\phi\|^2 \equiv \int d^4x \sqrt{g} (\delta\phi(x))^2 = \int d^4x \sqrt{\hat{g}} e^{2\phi(x)} (\delta\phi(x))^2.$$

Second, in order to define a short distance cutoff one should also use the metric $\hat{g}e^\phi$ itself: the cutoff fluctuates with the field. Let us follow David, Distler and Kawai [2],[3] and assume that the ϕ -dependence of the measure in (4.1) can be absorbed in a local renormalizable action:

$$D_{\hat{g}e^\phi} \phi e^{-S_i[\hat{g},\phi]} = D_{\hat{g}} \phi e^{-S_{loc}[\hat{g},\phi]}, \quad (4.2)$$

where now on the right-hand side

$$\|\delta\phi\|^2 \equiv \int d^4x \sqrt{\hat{g}} (\delta\phi(x))^2$$

and the cutoff no longer fluctuates.

What is S_{loc} ? Although the ϕ -dependence of the measure in (4.1) looks inconvenient we do learn something important from (4.1): simultaneously changing

$$\hat{g}_{\mu\nu} \rightarrow \hat{g}_{\mu\nu} e^{\phi_0}, \quad \phi \rightarrow \phi - \phi_0$$

does not change the measure or $S_{R^2}, S_R, S_{c.c.}$. It does change the induced action.

From (3.7) we see (reinstating the ϕ - independent terms (3.6) into (4.1)):

$$S_i[\hat{g}, \phi] \rightarrow S_i[\hat{g}, \phi] - S_i[\hat{g}, \phi_0]$$

We conclude

$$Z[\hat{g}e^{\phi_0}] = Z[\hat{g}] e^{S_i[\hat{g}, \phi_0]} \quad (4.3)$$

that is, the ϕ -theory behaves as if it were a conformal field theory with conformal anomaly (3.2) given by $a = -A_0$, $b = -B_0$. This is, of course, precisely what is needed in order to insure that the background metric is really a fake: if we vary it, $\hat{g}_{\mu\nu} \rightarrow \hat{g}_{\mu\nu}e^{\phi_0}$, the variation of the determinants in (3.10) is determined by their total conformal anomalies $+A_0$, $+B_0$, defined in (3.9), and that just cancels the $-A_0$, $-B_0$ from the ϕ theory.

So let us replace (4.1) as in (4.2) by a four dimensional conformal field theory with conformal anomaly given by

$$a = -a_O - a_L - a_{mat}, \quad b = -b_O - b_L - b_{mat}. \quad (4.4)$$

I will propose – and justify in a moment – that as in two dimensions S_{loc} in (4.2) is again the induced action with modified coefficients A, B and modified interactions:

$$Z[\hat{g}] \sim \int D_{\hat{g}} \phi \, e^{\frac{B}{32\pi^2} S_0[\hat{g}, \phi] + \frac{A}{32\pi^2} S_1[\hat{g}, \phi] - \eta_2 \hat{S}_{R^2} - \gamma_2 \hat{S}_R - \lambda_2 \hat{S}_{c.c.}} \quad (4.5)$$

where $\hat{S}_{R^2}$, $\hat{S}_R$, and $\hat{S}_{c.c.}$ are marginal operators of the free theory given by S_0 and S_1 and will be discussed below.

The free theory ($\eta_2, \gamma_2, \lambda_2 = 0$) of (4.5) has conformal anomaly

$$a = -A + a_M, \quad b = -B + b_M \quad (4.6)$$

where M is the operator (3.3). This can be seen as follows: Setting $\hat{g}_{\mu\nu} = \tilde{g}_{\mu\nu}e^{\phi_0}$ we

see from (3.8):

$$\begin{aligned}
& \int D_{\hat{g}} \phi \, e^{\frac{B}{32\pi^2} S_0[\hat{g}, \phi] + \frac{A}{32\pi^2} S_1[\hat{g}, \phi]} \\
&= \int D_{\hat{g}} \phi \, e^{\int d^4x \sqrt{\hat{g}} \left\{ \frac{B}{64\pi^2} [(\phi + \phi_0) \tilde{M}(\phi + \phi_0) - \phi_0 \tilde{M} \phi_0] + \frac{A}{32\pi^2} [\tilde{F}(\phi + \phi_0) - \tilde{F} \phi_0] \right\}} \quad (4.7) \\
&= e^{-\frac{B}{32\pi^2} S_0[\tilde{g}, \phi_0] - \frac{A}{32\pi^2} S_1[\tilde{g}, \phi_0]} \int D_{\hat{g}} \phi \, e^{\int d^4x (\sqrt{\hat{g}} \frac{B}{64\pi^2} \phi \hat{M} \phi + \frac{A}{32\pi^2} \hat{F} \phi)}
\end{aligned}$$

by shifting $\phi \rightarrow \phi + \phi_0$ and using the fact that $\sqrt{g}M$ and $\sqrt{g}F$ are conformally invariant. So $-A, -B$ are the “classical” contributions^{*} to (4.6) and a_M, b_M are the quantum contributions from M . Therefore we see from (4.4) that the ansatz (4.5) is consistent if

$$A = a_O + a_L + a_{mat} + a_M, \quad B = b_O + b_L + b_{mat} + b_M. \quad (4.8)$$

How do we know that a_M, b_M do not depend on the moduli m_i ? The only local scale invariant quantity they could depend on is $\int \sqrt{\hat{g}} \hat{F}$, which is a topological invariant in the case of $W_+ = 0$.

Why does the free part of S_{loc} in (4.2) have to be of the form of the free part of the induced action S_i again? One can plausibly, though not rigorously, argue as follows: there are two ways to obtain the right effective action (4.3) via (4.2); (a), S_{loc} is classically conformally invariant and a, b come purely from the quantum anomaly or (b), the “classical” variation of S_{loc} is of the form of the induced action S_i . In case (a) a and b would be just numbers that will in general not cancel the anomalies as needed in (4.4) (multiplying S_{loc} by a factor would then not change the conformal anomaly). Only in case (b) there are parameters like A, B in a, b that can be adjusted to satisfy (4.4). But the only local free action whose “classical” variation is the induced action is the induced action itself.

^{*} Here and below I will call these contributions also “anomalies,” although they actually arise from the fact that the action is classically not quite conformally invariant.

Let us now turn to the operators $\hat{S}_{R^2}$, $\hat{S}_R$ and $\hat{S}_{c.c.}$ in (4.5). The consistency condition of invariance under rescaling of the background metric (in particular, the theory is at a renormalization group fixed point) means that the integrands of $\hat{S}_{R^2}$, the “dressed” Hilbert-Einstein action $\hat{S}_R$ and the “dressed” cosmological constant $\hat{S}_{c.c.}$ must be locally scale invariant operators. Let us try the ansatz

$$\begin{aligned}\hat{S}_{c.c.} &= \int d^4x \sqrt{\hat{g}} e^{2\alpha\phi} \\ \hat{S}_R &= \int d^4x \sqrt{\hat{g}} e^{\beta\phi} (\hat{\nabla}\phi)^2 + \dots \\ \hat{S}_{R^2} &= \int d^4x \sqrt{\hat{g}} (\hat{\nabla}\phi)^4 + \dots\end{aligned}\tag{4.9}$$

with α, β , and “...” determined so that the integrands of (4.9) are scaling operators of conformal dimension 4, to cancel the -4 from $\sqrt{\hat{g}}$. In the language of string theory, they are vertex operators of our theory of noncritical three branes. All of them should be moduli deformations, if the background metric $\hat{g}$ is really fictitious. So far I have verified this only for $\hat{S}_{c.c.}$. The “...” includes possible corrections of order $\eta_2, \gamma_2, \lambda_2$ that may be needed in order to keep the other operators marginal as we move away from $\eta_2, \gamma_2, \lambda_2 = 0$. Some calculations with $\hat{S}_{R^2}, \hat{S}_R$ and $\hat{S}_{c.c.}$ can also be found in [7] (however $\alpha = \beta$ there).

To calculate the (classical plus anomalous) dimension of $e^{2\alpha\phi}$ with action (4.5) at $\eta_2, \gamma_2, \lambda_2 = 0$ one may go to conformally flat $\hat{g}_{\mu\nu} = e^{\phi_0} \delta_{\mu\nu}$ where $\hat{M} = 2 \hat{\square}^2$ and $S_1 = 0$. Because of the shift $\phi + \phi_0 \rightarrow \phi$ in (4.7), the condition

$$\dim(e^{2\alpha\phi}) = 4 \text{ with action } \sim S_0$$

is equivalent to the condition

$$\dim(e^{2\alpha\phi}) = 4 - 4\alpha \text{ with action } \sim \int d^4x \phi \square^2 \phi.\tag{4.10}$$

Due to the quartic propagator, this four dimensional theory is formally very similar to an ordinary free scalar field theory in two dimensions. In particular, $: e^{2\alpha\phi} :$ will

be a scaling operator. Its dimension in (4.10) is now purely anomalous. It is found from the two-point function

$$\langle e^{2\alpha\phi(x)} e^{-2\alpha\phi(y)} \rangle \sim e^{-4\alpha^2\Delta(|x-y|)} \sim |x-y|^{-\frac{8\alpha^2}{B}}, \quad (4.11)$$

where the propagator

$$\Delta(r) = \frac{2}{B} \log r \quad \text{with} \quad -\frac{B}{16\pi^2} \square^2 \Delta(r) = \delta(r)$$

of the free theory has been used. Thus, $\dim(e^{2\alpha\phi}) = \frac{4\alpha^2}{B}$, and (4.10) becomes:

$$4 - 4\alpha = \frac{4\alpha^2}{B}. \quad (4.12)$$

This determines α once B is known^{*} Similarly, β in $\hat{S}_R = \int \sqrt{\hat{g}} e^{\beta\phi} (\hat{\nabla}\phi)^2 + \dots$ is determined by requiring $e^{\beta\phi}$ to have dimension 2:

$$2 - 2\beta = \frac{\beta^2}{B}. \quad (4.13)$$

α and β are independent of the moduli m_i , for the same reason as a_M, b_M are.

The result for the dimension of the operator $e^{p\phi}$ agrees with the result of [7]. (Q^2 of [7] is $-2B$) Let me briefly point out two differences of section 4 with [7], where the theory of the conformal factor was studied as a ‘minisuperspace’ theory rather than as gravity with a self-duality constraint: First, I did not use the symmetry argument that was used in [7] to justify (for conformally flat $\hat{g}$ only) that S_{loc} in (4.2) is again the Liouville action. Second, in [7] α was equal to β in (4.9) with the consequence that the simultaneous presence of $\hat{S}_R$ and $\hat{S}_{c.c.}$ at the fixed point was inconsistent. This led to a suggestion about the cosmological constant problem, to a different value for α and will result in a different value for the analog of the ‘ $c = 1$ barrier’ of two dimensional gravity.

^{*} see the note added at the end of this paper.

As in two dimensions, if Φ_i is a scaling operator of the matter theory with conformal dimension Δ_i , the operator

$$O_i \equiv \int d^4x \sqrt{\hat{g}} e^{\gamma_i \phi} \Phi_i$$

with γ_i determined analogously to (4.12) by

$$4 - 2\gamma_i = \frac{\gamma_i^2}{B} + \Delta_i \quad (4.14)$$

is a marginal operator that can be added to the action, at least infinitesimally.

Provided that truly marginal operators $\hat{S}_{R^2}, \hat{S}_R$ can also be found, we can now rewrite (3.10) as

$$\begin{aligned} & \int \prod_i dm_i \det(O^\dagger O)_{\hat{g}}^{-\frac{1}{2}} \det(L^\dagger L)_{\hat{g}}^{\frac{1}{2}} \int D_{\hat{g}}x D_{\hat{g}}\phi e^{-S_{mat}[\hat{g},x] - S[\hat{g},\phi]}, \\ & S[\hat{g},\phi] = \frac{-B}{32\pi^2} S_0[\hat{g},\phi] + \frac{-A}{32\pi^2} S_1[\hat{g},\phi] \\ & \quad + \eta \hat{S}_{R^2}[\hat{g},\phi] + \gamma \hat{S}_R[\hat{g},\phi] + \lambda \hat{S}_{c.c.}[\hat{g},\phi]. \end{aligned} \quad (4.15)$$

The index of η, γ, λ has been dropped. (4.15) describes free fields plus marginal interactions in a gravitational instanton background. A, B are given by (4.8) and $S_0, S_1, \hat{S}_{R^2}, \hat{S}_R, \hat{S}_{c.c.}$ by (3.5) and (4.9).

5. Scaling

As an application of the preceding let us derive scaling laws by studying the integral over the constant mode of ϕ , as is done in two dimensions^{[3], [15]}. The fixed volume partition function at the critical point $\eta, \gamma, \lambda \sim 0$ is defined as

$$Z(V) \equiv \int \prod_i dm_i \chi(m_i) \int D_{\hat{g}} \phi e^{\frac{B}{32\pi^2} S_0[\hat{g}, \phi] + \frac{A}{32\pi^2} S_1[\hat{g}, \phi]} \delta\left(\int \sqrt{\hat{g}} e^{2\alpha\phi} - V\right) \quad (5.1)$$

with $\chi(m_i)$ as in (3.10). Under the constant shift $\phi \rightarrow \phi + c$ we see from (3.5):

$$\begin{aligned} \delta S_0 &= c \int d^4x \sqrt{\hat{g}} \hat{G} = 32\pi^2 c \chi \\ \delta S_1 &= c \int d^4x \sqrt{\hat{g}} \hat{F} = -48\pi^2 c \tau, \end{aligned}$$

where the topological invariants χ and τ are the Euler characteristic and signature of the manifold ($W_+ = 0$ here):^[16]

$$\tau = \frac{1}{48\pi^2} \int d^4x \sqrt{g} (W_+^2 - W_-^2) \quad \text{and} \quad \chi = \frac{1}{32\pi^2} \int d^4x \sqrt{g} G.$$

From this it follows that

$$\begin{aligned} Z(V) &= e^{(-2\alpha + B\chi - \frac{3}{2}A\tau)c} Z(e^{-2\alpha c} V) \\ \rightarrow Z(V) &\sim V^{-1 + \frac{1}{4\alpha}(2B\chi - 3A\tau)}. \end{aligned} \quad (5.2)$$

α is given in terms of B by (4.12). For the four sphere ($\chi = 2, \tau = 0$),

$$Z(V) \sim V^{-1 + \frac{B}{\alpha}}.$$

This quantity should be the easiest one to check with computer simulations.*

* See note added at the end of the paper for the value of the exponent

Inserting operators into (5.1) yields

$$< O_1 \dots O_n > (V) \sim V^{-1 + \frac{1}{4\alpha}(2B\chi - 3A\tau + 2\sum \gamma_i)}$$

with γ_i determined by (4.14). For nonzero cosmological constant λ one finds from

$$< O_1 \dots O_n >_\lambda = \int dV e^{-\lambda V} < O_1 \dots O_n >_0 (V) \quad (5.3)$$

the scaling behavior

$$< O_1 \dots O_n >_\lambda \sim \lambda^{-\frac{1}{4\alpha}(2B\chi - 3A\tau + 2\sum \gamma_i)}, \quad (5.4)$$

provided the integral (5.3) converges, i.e. $\frac{1}{4\alpha}(2B\chi - 3A\tau + 2\sum \gamma_i) > 0$. Otherwise there will be additional cutoff-dependent terms in (5.4). ^[15]

Replacing in (5.1)

$$\delta\left(\int \sqrt{\hat{g}} e^{2\alpha\phi} - V\right) \rightarrow \delta\left(\frac{\int \sqrt{\hat{g}} e^{\beta\phi} [(\nabla\phi)^2 + \dots]}{\int \sqrt{\hat{g}} e^{2\alpha\phi}} - \bar{R}\right)$$

one obtains the partition function for fixed curvature per volume at $\eta, \gamma, \lambda = 0$:

$$Z(\bar{R}) \sim \bar{R}^{-1 + \frac{B\chi}{\beta - 2\alpha} - \frac{3}{2} \frac{A\tau}{\beta - 2\alpha}}. \quad (5.5)$$

Scaling laws (5.2) and (5.4) are similar to the two dimensional ones.

6. Outlook

Fixed Points Of Gravity With A Weyl Term

Conformally self-dual gravity can also be understood as quantum gravity with the action

$$\int_M d^4x \sqrt{g} (\lambda + \gamma R + \eta R^2 + \rho W_+^2) \quad (6.1)$$

in the limit

$$\rho \rightarrow \infty. \quad (6.2)$$

(6.1) is the most general local renormalizable fourth order derivative action of four dimensional gravity, up to the topological invariants τ and χ of the previous section. In section 2, the metric was split into ϕ , diffeomorphisms ξ , moduli m_i and five $\bar{h}$ components. (6.2) can be understood as the “classical limit” for the $\bar{h}$ components, in which only the linearized W_+ -term $O^\dagger \bar{h}$ of (2.5) is important for the $\bar{h}$ -integral. This Gaussian integral can be performed at each point $g_0(\phi, \xi, m_i)$,

$$\int D\bar{h} e^{-\rho \int d^4x \sqrt{g} W_+^2} \sim \det(\rho O O^\dagger)_{g_0}^{-\frac{1}{2}} = \det(\rho O^\dagger O)_{\hat{g}e^\phi}^{-\frac{1}{2}}.$$

This leads again to the integral (2.6), our starting point. (The extra factor ρ only renormalizes the cosmological constant and does not influence the anomaly coefficients of section 3.)

Being fourth order in derivatives, the R^2 - and W_+^2 -terms will give rise to negative norm states. But this will not bother us in the limit $\rho \rightarrow \infty$, because the W_+^2 -term decouples and we can again tune away the R^2 -term in the end. Of course, the new fourth order derivative action S_0 is induced by the conformal anomaly as in section 3.

One might worry that the renormalization group flow will take us from $\rho \sim \infty$ to finite ρ so that the limit (6.2) does not make sense as an effective theory. However, since at $\rho \sim \infty$ the five $\bar{h}$ components decouple from the other five components of the

metric, $\rho \sim \infty$ corresponds to a renormalization group fixed point. More precisely, defining $\epsilon \equiv \frac{1}{\sqrt{\rho}}$, rescaling $\bar{h} \rightarrow \epsilon \bar{h}$ and expanding the action in ϵ , one obtains:

$$L_0[\phi, x] + \bar{h} O O^\dagger \bar{h} + \epsilon L_i[\bar{h}, \phi, x] + o(\epsilon^2) \quad (6.3)$$

where L_0 is the $\bar{h}$ -independent part, and L_i are interaction terms of $\bar{h}$ with itself, ϕ and x , x representing the matter. Thus the beta function for $\epsilon \sim \rho^{-\frac{1}{2}}$ will receive contributions only from diagrams that couple $\bar{h}$ and ϕ , so it will be at least of order ϵ and vanish as $\epsilon \rightarrow 0$. If $(\bar{\lambda}, \bar{\gamma}, \bar{\eta})$ is a fixed point of L_0 , then $(\bar{\lambda}, \bar{\gamma}, \bar{\eta}, \rho = \infty)$ will be a fixed point of (6.1).

Presently I do not know whether it will be infrared stable or unstable. If it turns out to be infrared stable, (1.1) describes a “conformally self-dual phase” of higher order derivative gravity – obviously not the world we live in. If it turns out to be ultraviolet stable, and if gravity with a Weyl term has something to do with reality, one could speculate that (1.1) describes gravity at short distances.

Hopefully the results found in section 5 will be the starting point for finding similar results for ρ finite or zero.

Topological gravity:

In two dimensional quantum gravity the correlation functions of local operators are related to the correlation functions of topological gravity^[10] which are intersection numbers of submanifolds on the moduli space of Riemann surfaces with punctures. Given the similarity of conformally self-dual quantum gravity to two dimensional quantum gravity, it would be very interesting to see if there is a similar relation between it and four dimensional topological gravity^[9]. This is suggested by the fact that the moduli space of the latter theory seems to be precisely the moduli space of conformally self-dual metrics that arose here. One could look for a matter system, analogous to the $c = -2$ system in two dimensions^[17] that, coupled to gravity, reproduces the BRST multiplet of 4D topological gravity. The relation between the “topological” and the “physical” phase of quantum gravity could then be studied in a less trivial model than the two dimensional one.

Conclusion

Surprisingly enough, methods of two dimensional quantum gravity can be applied to four dimensional quantum gravity at least in the limit of infinite Weyl coupling. The scaling laws are similar to those of two dimensions and can hopefully be compared with numerical simulations based on random triangulations.* It remains to be seen whether analogs of the Virasoro constraints decouple the negative norm states arising from the fourth order derivative term that is induced by the conformal anomaly.

Many other interesting questions could now be asked, but this will be left for the future.

Acknowledgements

I would like to thank K. Li, E. Mottola and M. Staudacher for discussions, J. H. Schwarz for advice and encouragement and E. Witten for suggesting to study conformally self-dual gravity.

Note added

The original version of this paper did not contain the values of a_O, b_O, a_L, b_L, a_M , and b_M . After it appeared, it was pointed out^{[18][19]} that the coefficients $a_O + a_L$ and $b_O + b_L$ have been computed long ago^[20] and that a_M, b_M for M in (3.3) can also be inferred from the literature^[21] with the results:

$$a_O + a_L = \frac{796}{120} \quad b_O + b_L = -\frac{1566}{360} \quad a_M = \frac{-8}{120} \quad b_M = -\frac{28}{360}, \quad (A.1)$$

see [18] for independent checks. We can now make some numerical predictions: with these results one obtains from (3.11) and (4.8)

$$A = \frac{1}{120}(N_0 + 6N_{\frac{1}{2}} + 12N_1 + 788), \quad B = -\frac{1}{360}(N_0 + 11N_{\frac{1}{2}} + 62N_1 + 1538) \quad (A.2)$$

where $N_0, N_{\frac{1}{2}}, N_1$ are the number of conformally coupled scalars, spin $\frac{1}{2}$ fermions and

* predicted values of exponents and analog of $c = 1$ barrier are given in the note below

massless gauge fields. (4.12), (4.13) and (4.14) become

$$\begin{aligned} 2\alpha &= -B - \sqrt{B^2 + 4B}, \\ \beta &= -B - \sqrt{B^2 + 2B}, \\ \gamma_i &= -B - \sqrt{B^2 + (4 - \Delta_i)B} \end{aligned} \tag{A.3}$$

Thus α will be real if $B \geq 0$ or $B \leq -4$. The second constraint is the relevant one since B is negative. The reality constraint $B \leq \Delta_i - 4$ on γ_i is weaker than the one for α in (A.3) as long as we allow only operators with positive dimension Δ_i . The signs in front of the square roots have been picked to give the correct results $\alpha = \beta = 1, \gamma_i = 2 - \frac{\Delta_i}{2}$ in the classical limit $B \rightarrow -\infty$.

To compare with two dimensional gravity, redefine the anomaly coefficient b of (3.2) as $\tilde{c} \equiv -360 b$, so that $\tilde{c}_0, \tilde{c}_{\frac{1}{2}}, \tilde{c}_1$ are 1, 11, and 62. $B \leq -4$ becomes

$$\tilde{c}_{mat} + \tilde{c}_L + \tilde{c}_O + \tilde{c}_M \geq 1440 \rightarrow \tilde{c}_{mat} \geq -98. \tag{A.4}$$

The analogous restriction in two dimensions is $c_{mat} \leq 1$, where c_{mat} is the matter central charge. If the cosmological constant term is absent, the barrier for $\tilde{c}$, rather than being -98 , is determined by the lowest dimension operator. We see that in pure gravity α is real. In contrast with two dimensions, the situation is improved by adding conventional matter like conformally coupled scalar fields, families of fermions or gauge fields. The $\tilde{c} = -98$ barrier would only be crossed by adding exotic matter with positive anomaly coefficient b in (3.2). It would be very interesting to investigate if the barrier becomes positive as we move away from $\rho \sim \infty$ in (6.1).

Using the values (A.1), we conclude that for conventional matter, on the sphere and at the critical point, $Z(V)$ always diverges (faster than V^{-1}) at small volumes and $Z(\bar{R})$ at large curvature per volume. E.g, for pure gravity on the sphere, (5.2), (5.5), (A.2) and (A.3) lead to the predictions:

$$Z(V) \sim V^{-3.675..} \qquad Z(\bar{R}) \sim \bar{R}^{+3.194..} \tag{A.5}$$

REFERENCES

1. A.M. Polyakov, Phys. Lett. 103B, 207 (1981)
2. F. David, Mod. Phys. Lett. A3, 1651 (1988)
3. J. Distler and H. Kawai, Nucl. Phys. B 321, 509 (1988)
4. V. Knizhnik, A. M. Polyakov and A. B. Zamolodchikov, Mod. Phys. Lett. A3, 819 (1988)
5. M. E. Agishtein and A. A. Migdal, Nucl. Phys. B350, 690 (1991)
6. E.g., D. J. Gross and A. A. Migdal, Nucl. Phys. B 340, 333 (1990) and references therein.
7. I. Antoniadis and E. Mottola, Los Alamos preprint LA-UR-91-1653 (1991)
8. E. D' Hoker, Mod. Phys. Lett. A6, 745 (1991)
9. E. Witten, Phys. Lett. B 206, 601 (1988); J.M.F. Labastida and M. Pernici, Phys. Lett. B 212, 56 (1988)
10. E. Witten, Nucl. Phys. B 340, 281 (1990); E. and H. Verlinde, Nucl. Phys. B 348, 457 (1991)
11. L. Alvarez-Gaumé and E. Witten, Nucl. Phys. B 234, 269 (1983)
12. In T. Eguchi, P. B. Gilkey and A. J. Hanson, Phys. Rep. 66, 213 (1988), after I. M. Singer
13. E.g., M. J. Duff, Nucl. Phys. B125, 334 (1977); N. D. Birrell and P. C. W. Davies, "Quantum Fields in Curved Space," Cambridge University Press 1982
14. E.g., M.B. Green, J.H. Schwarz and E. Witten, "Superstring Theory", Cambridge University Press 1987
15. N. Seiberg, Prog. Theor. Phys. S 102, 319 (1990)
16. E.g., M.F. Atiyah, N.J. Hitchin and I.M. Singer, Proc. Roy. Soc. London A362,425 (1978)

17. J. Distler, Nucl. Phys. B342, 523 (1990)
18. I. Antoniadis, P.O. Mazur and E. Mottola, preprint CPTH-A173.0492 / LA-UR-92-1483 (1992)
19. E. Elizalde and S.D. Odintsov, Barcelona preprint UB-ECM-PF 92/12 (1992)
20. E.S. Fradkin and A.A. Tseytlin, Nucl. Phys B 201, 469 (1982)
21. A.O. Barvinsky and G.A. Vilkovisky, Phys. Rep. 119, 1 (1985)